


SAFETY NOTICE on Covid-19 Vaccination

24.05.2021

- 
- **Vaccination can be done** in the local medical institutions as well as in some private clinics, such as Inter teach LLP and Mediker LLP. In some cases, mobile vaccination posts are arranged (e.g. in the territory of the shopping centers) or mobile vaccination teams.
 - **Vaccination is done in two stages:** at first with component I, then in **45 days** excluding the day of vaccination with component II. If it is not possible to get vaccinated with component II in 45 days, the vaccination can be done within 90 days.
 - **The persons who completed the vaccination are issued with a vaccination passport.** The passport is available via the eGov Mobile.
 - **The first immune response** is being generated at the 12th -14th day after the injection of component I, but the stable immunity will be formed not earlier than in **32-56 days**.

- Before vaccination, **interview and examination** by a physician is carried out, including measurement of body temperature, saturation, heart rate, blood pressure, auscultation of the respiratory and cardiovascular system, and pharyngeal examination.
- All vaccinated persons will spend the first 30 minutes **under medical supervision** in a medical facility. **For the next three days**, they will be under **the care of local physicians at the place of residence**.
- Each vaccinated person will be given a special **safety notice** indicating possible side effects of the vaccine and telephone numbers of clinics to contact in case of such effects



Possible Reaction to the Vaccination

On the first or second day after vaccination, the following symptoms may develop and shall disappear within the next **three days**:

- short flu-like syndrome (chills, fever, general feeling of being unwell, headache),
- local reactions (soreness at the injection area, hyperemia, swelling),
- less common are nausea, loss of appetite,
- allergic reactions may occur as well.

The vaccination **IS PROHIBITED** for the following persons:

- Below 18 years old.
- Those who have temporary or permanent medical restrictions against any vaccination.
- Those who have hypersensitivity to any component of the vaccine or a vaccine containing similar components.
- Those who had severe allergic reactions anamnesis (previously during one's life).
- Pregnant and breast feeding.
- **Contraindications against injection of the II component:** severe complications after the vaccination with the I component (anaphylactic shock, severe allergic reaction, convulsive attack, body temperature reaching 40°C).



Contraindications and Restrictions

The vaccination **SHALL BE POSTPONED**, if:

- Your body temperature is or above 37°C .
- You are suffering from **the acute** infectious or noninfectious disease (the vaccination shall be carried out in **2-4 weeks** after the recovery).
- You are suffering from the worsening of the chronic disease (the vaccination shall be carried out in **2-4 weeks** after the recovery or remission).
- You were in contact with COVID-19 case (vaccination shall be carried out **after the quarantine is over**).
- **You have recovered form COVID-19** (vaccination shall be carried out in **6 months** after the recovery taking into account the health condition before the vaccination).
- You were drinking alcohol within the last three days.

The necessity to be vaccinated shall be **CAREFULLY CONSIDERED** in case you have been diagnosed with:

- a chronic disease of liver and kidneys;
- diabetes mellitus;
- autoimmune diseases;
- lungs disease, asthma;
- severe diseases of hematogenesis system;
- epilepsy, cerebrovascular accident, and other disease of the central nerve system;
- diseases of cardiovascular system;
- allergy, atopy or eczema;
- malignant neoplasms.





HSE
HEALTH SAFETY ENVIRONMENT